

## **Outcome Reports**

The outcomes shown below have been collected using our state-of-the-art outcome system, the Astym Analyst™. Outcomes can be reported by six different outcome criteria including pain, functional status, pain and function combined functional improvement, patient satisfaction, and patient goals. In addition, data can be analyzed and displayed by the following factors:

• Age

Limitations on Visits

Percentage of Astym Tx. Visits

Chronicity

Mechanism of Onset

Prior Therapy Received

Comorbidities

Other Therapy Received

Referral Source

Gender

Patient Classification

Industry Code

The following are a few examples of report data available in the Analyst. If you would like to view reports in addition to those displayed below, please contact Performance Dynamics, Inc. at 765-751-2341.

## Functional Improvement in Work/Sport/Activity

Displayed by Mechanism of Onset

By Mechanism of Onset	Total Cases	Avg # of Astym Tx	Improved	Same	Worse
Minor trauma (strains, sprains, etc)	1972	8	93.9%	5.8%	0.4%
Major trauma (MVA, etc)	586	9	92.5%	6.8%	0.7%
Post-surgical	826	9	92.9%	6.4%	0.7%
Repetitive stress (specific activity)	3393	9	92.1%	7.4%	0.5%
Insidious onset (no specific activity)	3266	9	92.0%	7.6%	0.4%
Degenerative (joint)	186	8	91.4%	8.6%	0.0%
TOTALS	10,229	9	92.5%	7.1%	0.5%

<sup>\*</sup>Cases with no impairment IE and DC for functional improvement not included

## **Pain Ratings**

Displayed by Chronicity

By Chronicity	Total Cases	Avg # of Astym Tx	Excellent/Good	Fair/Poor	
Acute (0-14 days)	492	7	96.5%	3.5%	
Subacute (15 days to 12 weeks)	3344	8	92.8%	7.2%	
Chronic long term (> 12 weeks)	6492	9	87.9%	12.1%	
TOTALS	10,328	9	92.4%	7.6%	

<sup>\*</sup>Calculated from % in pain and D/C pain

## **Patient Satisfaction**

Displayed by Patient Classification

By Patient Classification	Total Cases	Avg # of Astym Tx	Satisfied	Neither	Unsatisfied
Recreationally active	3820	9	90.1%	8.4%	1.5%
Sedentary	1499	8	87.9%	9.3%	2.7%
Competitive athlete	1118	8	93.8%	5.1%	1.1%
Recreational athlete	1893	9	93.2%	5.7%	1.1%
Workers compensation	1708	9	85.6%	12.4%	2.0%
Non-workers compensation	290	8	89.7%	9.3%	1.0%
TOTALS	10,328	9	90.1%	8.4%	1.6%

Astym® Therapy Outcome Report Functional Improvement in Work/Sport/Activity

Functional Improvement in Work/Sport/Activity						
All Diagnostic Categories	Total Cases	Avg # of	Improved	Same	Worse	
Shoulder Girdle		Astym Tx	-			
Adhesive capsulitis	64	9	96.9%	1.6%	1.6%	
Biceps tendinopathy	75	8	85.4%	13.3%	1.3%	
Rotator cuff-impingement/tendinopathy	234	7	94.0%	4.7%	1.3%	
Rotator cuff-partial tear	40	8	75.0%	20.0%	2.5%	
Scapular dysfunction	40	7	90.0%	10.0%	0.0%	
Shoulder pain	215	7	89.8%	9.8%	0.5%	
Subacromial bursitis	30	7	96.7%	3.3%	0.0%	
Surgical repair of rotator cuff tear	38	8	89.5%	10.5%	0.0%	
Elbow/Forearm Cubital tunnel syndrome	29	9	86.2%	13.8%	0.0%	
Elbow pain	102	8	93.1%	6.9%	0.0%	
Lateral epicondylopathy	1480	10	91.2%	8.4%	0.5%	
Lateral epicondylopathy + carpal tunnel syndrome	70	9	88.5%	10.0%	1.4%	
Lateral epicondylopathy + medial epicondylopathy	160	10	88.7%	10.6%	0.6%	
Medial epicondylopathy	217	9	92.7%	5.5%	1.8%	
Radial tunnel syndrome	24	8	95.9%	4.2%	0.0%	
Other tendinopathy of forearm (sprain/strain forearm)	133	8	92.5%	7.5%	0.0%	
Ulnar neuritis	27	8	92.6%	7.4%	0.0%	
Hand & Wrist		I	T	T		
Carpal tunnel syndrome	295	9	87.8%	11.9%	0.3%	
DeQuervain's  Post surgical carpal tuppel release	174 37	8	87.9% 86.5%	11.5% 8.1%	0.6% 5.4%	
Post-surgical carpal tunnel release  Sprain-thumb/fingers	37	8	94.1%	5.9%	0.0%	
Sprain-unumb/lingers Sprain-wrist	134	7	94.1%	2.2%	0.0%	
Trigger finger	61	8	85.2%	13.1%	1.6%	
Fractured distal radius and ulna	25	11	100.0%	0.0%	0.0%	
Other Upper Extremity Diagnoses		•				
Scar management upper extremity-surgical or traumatic	49	9	98.0%	2.0%	0.0%	
2 or more diagnoses from elbow to fingers	160	9	89.4%	8.8%	1.9%	
Proximal and distal upper extremity pathologies	63	10	90.5%	6.3%	3.2%	
Hip & Groin		1	T .	1		
Groin sprain/strain	95	8	96.8%	3.2%	0.0%	
Hip pain	203	8	93.6%	6.4%	0.0%	
IT band syndrome/TFL Piriformis syndrome	111 49	8 7	92.8% 85.7%	7.2% 12.2%	0.0% 2.0%	
Trochanteric bursitis	157	8	98.1%	1.3%	0.6%	
Knee	107		30.170	1.070	0.070	
Hamstring strain	291	8	97.6%	2.4%	0.0%	
IT band syndrome	109	7	98.2%	1.8%	0.0%	
Knee DJD	35	8	94.3%	5.7%	0.0%	
Knee pain	289	7	93.0%	6.9%	0.0%	
Patellar tendinopathy	175	8	89.7%	10.3%	0.0%	
Patellofemoral pain, patellar instability, plica syndrome	181	8	90.1%	9.9%	0.0%	
Pes anserine bursitis	21	9	95.2%	4.8%	0.0%	
Post-surgical ACL/PCL reconstruction	44	9	93.1%	6.8%	0.0%	
Post-surgical arthroscopy  Post-surgical total knee arthroplasty	89 190	8 9	95.5% 96.3%	3.4% 3.7%	1.1% 0.0%	
Quad strain	69	6	94.2%	4.3%	1.4%	
Sprain-MCL/LCL	26	7	96.1%	3.8%	0.0%	
Shin, Ankle & Foot		· · · · · · · · · · · · · · · · · · ·			212.72	
Achilles tendinopathy	557	9	95.1%	4.7%	0.2%	
Ankle/foot pain	279	8	93.2%	6.1%	0.7%	
Calf strain	133	8	96.2%	3.8%	0.0%	
Plantar fasciopathy-heel pain	1445	9	91.2%	8.4%	0.3%	
Post-surgical Achilles tendon injury	42	10	95.2%	4.8%	0.0%	
Post-surgical hallux valgus/bunionectomy	33	9	100.0%	0.0%	0.0%	
Shin splints-tendinopathy of tibialis anterior, tibialis posterior, peroneal tendinopathy	194	9	96.4%	3.6%	0.0%	
Sprain-ankle Fractured ankle	123 40	8 10	98.4% 100.0%	1.6% 0.0%	0.0%	
Other Lower Extremity Diagnoses	40	10	100.0%	0.0%	0.0%	
Scar management lower extremity-surgical or traumatic	51	9	92.1%	7.8%	0.0%	
2 or more diagnoses from the knee to the toes	30	11	86.6%	13.3%	0.0%	
Proximal and distal lower extremity pathologies	26	8	92.3%	7.7%	0.0%	
Head & Neck						
Cervical discogenic pain	35	7	94.3%	5.7%	0.0%	
Cervical sprain/strain	69	7	94.2%	5.8%	0.0%	
Neck pain/cervicalgia	169	7	91.1%	8.9%	0.0%	
Torso/Rib	_	1		1		
Thoracic pain	63	8	90.5%	9.5%	0.0%	
Thoracic sprain/strain	23	6	95.6%	4.3%	0.0%	
Back Low back pain	101	7	96 70/	12.70/	0.69/	
Low back pain  Lumbar discogenic pain	181 25	6	86.7% 96.0%	12.7% 4.0%	0.6% 0.0%	
Sacroiliac sprain/strain	26	8	96.0%	3.8%	0.0%	
TOTALS	9688	9	92.0%	7.0%	0.0%	
		· -	J-1-70			

Outcomes above represent the following: Diagnostic Categories with more than 20 entries